

On a system of difference equations of third order solved in closed form

Youssef Akrouf   ^{1,4}, Nouressadat Touafek  ^{*2,4} and Yacine Halim  ^{**3,4}

¹Département des Sciences Exactes et d'Informatique, École Normale Supérieure Assia Djebar, Constantine, Algeria

²Department of Mathematics, Mohamed Seddik Ben Yahia University, Jijel, Algeria

³Department of Mathematics and Computer Science, Abdelhafid Boussouf University Center, Mila, Algeria

⁴LMAM Laboratory, Mohamed Seddik Ben Yahia University, Jijel, Algeria

Received 18 November 2021, Accepted 06 December 2021, Published 22 December 2021

Abstract. In this work, we show that the system of difference equations

$$x_{n+1} = \frac{ay_{n-2}x_{n-1}y_n + bx_{n-1}y_{n-2} + cy_{n-2} + d}{y_{n-2}x_{n-1}y_n},$$

$$y_{n+1} = \frac{ax_{n-2}y_{n-1}x_n + by_{n-1}x_{n-2} + cx_{n-2} + d}{x_{n-2}y_{n-1}x_n},$$

where $n \in \mathbb{N}_0$, x_{-2} , x_{-1} , x_0 , y_{-2} , y_{-1} and y_0 are arbitrary nonzero real numbers and a , b , c and d are arbitrary real numbers with $d \neq 0$, can be solved in a closed form.

We will see that when $a = b = c = d = 1$ the solutions are expressed using the famous Tetranacci numbers. In particular, the results obtained here extend those in our recent work.

Keywords: System of difference equations, general solution, Tetranacci numbers.

2020 Mathematics Subject Classification: 39A05, 39A06, 39A10.

1 Introduction

Nonlinear difference equations and their systems are hot topics that attract the attention of several researchers. A significant number of papers are devoted to this field of research. One can consult, for example, the papers [3, 5–18, 20–23, 26, 27, 30, 31, 36–44, 46], where one can find concrete models of such equations and systems, as well as understand the techniques used to solve them and investigate the behavior of their solutions. Recently, in [1] and as a

[✉] Corresponding author: youssef.akrouf@gmail.com

*ntouafek@gmail.com

**halyacine@yahoo.fr

generalization of the equations and systems studied in [4,19,32,45], we have solved in a closed form the system of difference equations

$$\begin{cases} x_{n+1} = \frac{ay_n x_{n-1} + bx_{n-1} + c}{x_{n-1} y_n}, \\ y_{n+1} = \frac{ax_n y_{n-1} + by_{n-1} + c}{y_{n-1} x_n}. \end{cases} \quad (1.1)$$

Here, and motivated by the above papers, one shows that one can express in closed form the well-defined solutions of the following system of difference equations

$$\begin{cases} x_{n+1} = \frac{ay_{n-2} x_{n-1} y_n + bx_{n-1} y_{n-2} + cy_{n-2} + d}{y_{n-2} x_{n-1} y_n}, \\ y_{n+1} = \frac{ax_{n-2} y_{n-1} x_n + by_{n-1} x_{n-2} + cx_{n-2} + d}{x_{n-2} y_{n-1} x_n}, \end{cases} \quad (1.2)$$

where $n \in \mathbb{N}_0$, the initial values x_{-2} , x_{-1} , x_0 , y_{-2} , y_{-1} and y_0 are arbitrary nonzero real numbers and the parameters a , b , c and d are arbitrary real numbers with $d \neq 0$.

Clearly if $d = 0$, then System (1.2) is nothing other than system (1.1). For the readers interested in the solutions of this system, one refers to [1], where the system (1.1) has been completely solved.

Noting also that the system (1.2) can be seen as a generalization of the equation

$$x_{n+1} = \frac{ax_{n-2} x_{n-1} x_n + bx_{n-1} x_{n-2} + cx_{n-2} + d}{x_{n-2} x_{n-1} x_n}, \quad n \in \mathbb{N}_0. \quad (1.3)$$

In fact, the solutions of (1.3) can be obtained from the solutions of (1.2) by choosing $y_{-i} = x_{-i}$, $i = 0, 1, 2$. The equation (1.3) was the subject of a substantial part of the paper [4], which also motivated our present study. The same equation was studied in complex numbers by Stevic in [29].

We will see that the explicit formulas of the well defined solutions of system (1.2) are expressed using the terms of the sequence $(J_n)_{n=0}^{+\infty}$ which are the solutions of the fourth-order linear homogeneous difference equation defined by the relation

$$J_{n+4} = aJ_{n+3} + bJ_{n+2} + cJ_{n+1} + dJ_n, \quad n \in \mathbb{N}_0, \quad (1.4)$$

and the special initial values

$$J_0 = 0, \quad J_1 = 0, \quad J_2 = 1 \text{ and } J_3 = a. \quad (1.5)$$

In this article one solves in closed form the equation (3.3). This well-known equation (with the same or different initial values and parameters) was the subject of some papers in the literature, see for example [25,29,47].

The characteristic equation associated to (3.3) is

$$\lambda^4 - a\lambda^3 - b\lambda^2 - c\lambda - d = 0, \quad (1.6)$$

and let α , β , γ and δ its four roots, then

$$\begin{cases} \alpha + \beta + \gamma + \delta = a \\ \alpha\beta + \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta + \gamma\delta = -b \\ \alpha\beta\gamma + \alpha\beta\delta + \alpha\gamma\delta + \beta\gamma\delta = c \\ \alpha\beta\gamma\delta = -d \end{cases} \quad (1.7)$$

One has:

Case 1: If all roots are real and equal. In this case,

$$J_n = (c_1 + c_2n + c_3n^2 + c_4n^3) \alpha^n.$$

Now using (1.7) and the fact that $J_0 = 0, J_1 = 0, J_2 = 1$ and $J_3 = a$, one obtains

$$J_n = \left(\frac{-n + n^3}{6\alpha^2} \right) \alpha^n. \quad (1.8)$$

Case 2: If three roots are real and equal, i.e. $\beta = \gamma = \delta$. In this case

$$J_n = c_1\alpha^n + (c_2 + c_3n + c_4n^2) \beta^n.$$

Now using (1.7) and the fact that $J_0 = 0, J_1 = 0, J_2 = 1$ and $J_3 = a$, one obtains

$$J_n = \frac{-\alpha}{(\beta - \alpha)^3} \alpha^n + \left(\frac{\alpha}{(\beta - \alpha)^3} - \frac{n(\alpha + \beta)}{2\beta(\beta - \alpha)^2} + \frac{n^2}{2\beta(\beta - \alpha)} \right) \beta^n, \quad (1.9)$$

Case 3: If two real roots are equal, i.e. $\gamma = \delta$. In this case

$$J_n = c_1\alpha^n + c_2\beta^n + (c_3 + c_4n) \gamma^n.$$

Now using (1.7) and the fact that $J_0 = 0, J_1 = 0, J_2 = 1$ and $J_3 = a$, one obtains

$$J_n = \frac{-\alpha}{(\gamma - \alpha)^2(\beta - \alpha)} \alpha^n + \frac{\beta}{(\gamma - \beta)^2(\beta - \alpha)} \beta^n + \left(\frac{\alpha\beta - \gamma^2}{(\gamma - \alpha)^2(\gamma - \beta)^2} + \frac{n}{(\gamma - \alpha)(\gamma - \beta)} \right) \gamma^n, \quad (1.10)$$

Case 4: If two double real roots are equal, i.e. $\alpha = \beta \neq \gamma = \delta$. In this case

$$J_n = (c_1 + c_2n) \alpha^n + (c_3 + c_4n) \gamma^n.$$

Now using (1.7) and the fact that $J_0 = 0, J_1 = 0, J_2 = 1$ and $J_3 = a$, one obtains

$$J_n = \left(\frac{\gamma + \alpha}{(\gamma - \alpha)^3} + \frac{n}{(\gamma - \alpha)^2} \right) \alpha^n + \left(-\frac{\gamma + \alpha}{(\gamma - \alpha)^3} + \frac{n}{(\gamma - \alpha)^2} \right) \gamma^n, \quad (1.11)$$

Case 5: If all the roots are real and different. In this case

$$J_n = c_1\alpha^n + c_2\beta^n + c_3\gamma^n + c_4\delta^n.$$

Again, using (1.7) and the fact that $J_0 = 0, J_1 = 0, J_2 = 1$ and $J_3 = a$, one obtains

$$J_n = \frac{-\alpha}{(\delta - \alpha)(\gamma - \alpha)(\beta - \alpha)} \alpha^n + \frac{\beta}{(\delta - \beta)(\gamma - \beta)(\beta - \alpha)} \beta^n + \frac{-\gamma}{(\delta - \gamma)(\gamma - \beta)(\gamma - \alpha)} \gamma^n + \frac{\delta}{(\delta - \gamma)(\delta - \beta)(\delta - \alpha)} \delta^n. \quad (1.12)$$

Case 6: If two real roots are equal, i.e. $\alpha = \beta$ and two roots are complex conjugate, i.e. $\delta = \bar{\gamma}$. In this case

$$J_n = (c_1 + c_2n)\alpha^n + c_3\gamma^n + c_4\bar{\gamma}^n.$$

Again, using (1.7) and the fact that $J_0 = 0$, $J_1 = 0$, $J_2 = 1$ and $J_3 = a$, one obtains

$$J_n = \left(\frac{\bar{\gamma}\gamma - \alpha^2}{(\bar{\gamma} - \alpha)^2(\gamma - \alpha)^2} + \frac{n}{(\bar{\gamma} - \alpha)(\gamma - \alpha)} \right) \alpha^n + \frac{-\gamma}{(\bar{\gamma} - \gamma)(\gamma - \alpha)^2} \gamma^n + \frac{\bar{\gamma}}{(\bar{\gamma} - \gamma)(\bar{\gamma} - \alpha)^2} \bar{\gamma}^n. \quad (1.13)$$

Case 7: If two real roots α , β are different and two roots are complex conjugate, i.e. $\delta = \bar{\gamma}$. In this case

$$J_n = c_1\alpha^n + c_2\beta^n + c_3\gamma^n + c_4\bar{\gamma}^n.$$

Again, using (1.7) and the fact that $J_0 = 0$, $J_1 = 0$, $J_2 = 1$ and $J_3 = a$, one obtains

$$J_n = \frac{-\alpha}{(\bar{\gamma} - \alpha)(\gamma - \alpha)(\beta - \alpha)} \alpha^n + \frac{\beta}{(\bar{\gamma} - \beta)(\gamma - \beta)(\beta - \alpha)} \beta^n + \frac{-\gamma}{(\bar{\gamma} - \gamma)(\gamma - \beta)(\gamma - \alpha)} \gamma^n + \frac{\bar{\gamma}}{(\bar{\gamma} - \gamma)(\bar{\gamma} - \beta)(\bar{\gamma} - \alpha)} \bar{\gamma}^n. \quad (1.14)$$

Case 8: If two complex roots are equal, i.e. $\alpha = \gamma$ and $\beta = \delta = \bar{\alpha}$. In this case

$$J_n = (c_1 + c_2n)\alpha^n + (c_3 + c_4n)\bar{\alpha}^n.$$

Again, using (1.7) and the fact that $J_0 = 0$, $J_1 = 0$, $J_2 = 1$ and $J_3 = a$, one obtains

$$J_n = \left(\frac{\bar{\alpha} + \alpha}{(\bar{\alpha} - \alpha)^3} + \frac{n}{(\bar{\alpha} - \alpha)^2} \right) \alpha^n + \left(\frac{-\bar{\alpha} - \alpha}{(\bar{\alpha} - \alpha)^3} + \frac{n}{(\bar{\alpha} - \alpha)^2} \right) \bar{\alpha}^n. \quad (1.15)$$

Case 9: If the roots are all complex and different, i.e. $\beta = \bar{\alpha}$ and $\delta = \bar{\gamma}$. In this case

$$J_n = c_1\alpha^n + c_2\bar{\alpha}^n + c_3\gamma^n + c_4\bar{\gamma}^n.$$

Again, using (1.7) and the fact that $J_0 = 0$, $J_1 = 0$, $J_2 = 1$ and $J_3 = a$, one obtains

$$J_n = \frac{-\alpha}{(\bar{\gamma} - \alpha)(\gamma - \alpha)(\bar{\alpha} - \alpha)} \alpha^n + \frac{\bar{\alpha}}{(\bar{\gamma} - \bar{\alpha})(\gamma - \bar{\alpha})(\bar{\alpha} - \alpha)} \bar{\alpha}^n + \frac{-\gamma}{(\bar{\gamma} - \gamma)(\gamma - \bar{\alpha})(\gamma - \alpha)} \gamma^n + \frac{\bar{\gamma}}{(\bar{\gamma} - \gamma)(\bar{\gamma} - \bar{\alpha})(\bar{\gamma} - \alpha)} \bar{\gamma}^n. \quad (1.16)$$

2 The main theorem and some particular cases

Here, one gives a closed form for the well defined solutions of the system (1.2) with $d \neq 0$. One will use the same change of variables as in [1] to transform the system (1.2) to a linear one and then follows the same procedure as in [1] to obtain the closed-form of the solutions. To get the solutions of the corresponding linear system, one needs to solve some fourth-order linear difference equations. In particular, one derives from the main result (Main Theorem), for which one leaves the proof to the next section, the solutions of some particular systems and equations where their solutions are related to the famous Tetranacci numbers.

One recalls that by a well defined solution of system (1.2), one means a solution that satisfies $x_n y_n \neq 0$, $n \geq -2$. The set of well defined solutions is not empty. In fact, it suffices to choose the initial values and the parameters a , b , c and d positive, to see that every solution of (1.2) will be well defined.

2.1 Closed form of well defined solutions of the system (1.2)

The following result gives an explicit formula for well-defined solutions of the system (1.2).

Theorem 2.1. (Main Theorem) Let $\{x_n, y_n\}_{n \geq -2}$ be a well defined solution of (1.2). Then, for $n \in \mathbb{N}_0$, one has

$$\begin{aligned} x_{2n+1} &= \frac{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1})y_{-2} + (J_{2n+4} - aJ_{2n+3})x_{-1}y_{-2} + J_{2n+3}y_0x_{-1}y_{-2}}{dJ_{2n+1} + (cJ_{2n+1} + dJ_{2n})y_{-2} + (J_{2n+3} - aJ_{2n+2})x_{-1}y_{-2} + J_{2n+2}y_0x_{-1}y_{-2}}, \\ x_{2n+2} &= \frac{dJ_{2n+3} + (cJ_{2n+3} + dJ_{2n+2})x_{-2} + (J_{2n+5} - aJ_{2n+4})y_{-1}x_{-2} + J_{2n+4}x_0y_{-1}x_{-2}}{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1})x_{-2} + (J_{2n+4} - aJ_{2n+3})y_{-1}x_{-2} + J_{2n+3}x_0y_{-1}x_{-2}}, \\ y_{2n+1} &= \frac{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1})x_{-2} + (J_{2n+4} - aJ_{2n+3})y_{-1}x_{-2} + J_{2n+3}x_0y_{-1}x_{-2}}{dJ_{2n+1} + (cJ_{2n+1} + dJ_{2n})x_{-2} + (J_{2n+3} - aJ_{2n+2})y_{-1}x_{-2} + J_{2n+2}x_0y_{-1}x_{-2}}, \\ y_{2n+2} &= \frac{dJ_{2n+3} + (cJ_{2n+3} + dJ_{2n+2})y_{-2} + (J_{2n+5} - aJ_{2n+4})x_{-1}y_{-2} + J_{2n+4}y_0x_{-1}y_{-2}}{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1})y_{-2} + (J_{2n+4} - aJ_{2n+3})x_{-1}y_{-2} + J_{2n+3}y_0x_{-1}y_{-2}} \end{aligned}$$

where the initial values $x_{-2}, x_{-1}, x_0, y_{-2}, y_{-1}$ and $y_0 \in (\mathbb{R} - \{0\}) - F$, with F is the Forbidden set of system (1.2) given by

$$F = \bigcup_{n=0}^{\infty} \{(x_{-2}, x_{-1}, x_0, y_{-2}, y_{-1}, y_0) \in (\mathbb{R} - \{0\}) : A_n = 0 \text{ or } B_n = 0\},$$

where

$$\begin{aligned} A_n &= dJ_{n+1} + (cJ_{n+1} + dJ_n)y_{-2} + (J_{n+3} - aJ_{n+2})x_{-1}y_{-2} + J_{n+2}y_0x_{-1}y_{-2}, \\ B_n &= dJ_{n+1} + (cJ_{n+1} + dJ_n)x_{-2} + (J_{n+3} - aJ_{n+2})y_{-1}x_{-2} + J_{n+2}x_0y_{-1}x_{-2}. \end{aligned}$$

2.2 Particular cases

Now, we focus our study on some particular cases of system (1.2).

2.2.1 The solutions of the equation $x_{n+1} = (ax_{n-2}x_{n-1}x_n + bx_{n-1}x_{n-2} + cx_{n-2} + d) / (x_{n-2}x_{n-1}x_n)$

If one chooses $y_{-2} = x_{-2}, y_{-1} = x_{-1}$ and $y_0 = x_0$, then system (1.2) is reduced to the equation

$$x_{n+1} = \frac{ax_{n-2}x_{n-1}x_n + bx_{n-1}x_{n-2} + cx_{n-2} + d}{x_{n-2}x_{n-1}x_n}, \quad n \in \mathbb{N}_0. \quad (2.1)$$

So, it follows from the **Main Theorem**

Corollary 2.2. Let $\{x_n\}_{n \geq -2}$ be a well defined solution of the equation (2.1). Then for $n \in \mathbb{N}_0$, one has

$$\begin{aligned} x_{2n+1} &= \frac{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1})x_{-2} + (J_{2n+4} - aJ_{2n+3})x_{-1}x_{-2} + J_{2n+3}x_0x_{-1}x_{-2}}{dJ_{2n+1} + (cJ_{2n+1} + dJ_{2n})x_{-2} + (J_{2n+3} - aJ_{2n+2})x_{-1}x_{-2} + J_{2n+2}x_0x_{-1}x_{-2}}, \\ x_{2n+2} &= \frac{dJ_{2n+3} + (cJ_{2n+3} + dJ_{2n+2})x_{-2} + (J_{2n+5} - aJ_{2n+4})x_{-1}x_{-2} + J_{2n+4}x_0x_{-1}x_{-2}}{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1})x_{-2} + (J_{2n+4} - aJ_{2n+3})x_{-1}x_{-2} + J_{2n+3}x_0x_{-1}x_{-2}}. \end{aligned}$$

It is worth noting that this equation was studied in [4, 29].

2.3 The solutions of the system (1.2) with $a = b = c = d = 1$

Consider the system

$$\begin{cases} x_{n+1} = \frac{y_{n-2}x_{n-1}y_n + x_{n-1}y_{n-2} + y_{n-2} + 1}{y_{n-2}x_{n-1}y_n}, \\ y_{n+1} = \frac{x_{n-2}y_{n-1}x_n + y_{n-1}x_{n-2} + x_{n-2} + 1}{x_{n-2}y_{n-1}x_n}, \end{cases} n \in \mathbb{N}_0, \quad (2.2)$$

which is a particular case of the system (1.2) with $a = b = c = d = 1$. In this case the sequence $\{J_n\}$ is nothing other than the sequence of Tetranacci numbers $\{T_n\}$, that is

$$T_{n+4} = T_{n+3} + T_{n+2} + T_{n+1} + T_n, \quad n \in \mathbb{N}_0, \quad \text{where } T_0 = T_1 = 0, T_2 = 1 \text{ and } T_3 = 1,$$

and one has

$$T_n = \frac{-\alpha}{(\bar{\gamma} - \alpha)(\gamma - \alpha)(\beta - \alpha)} \alpha^n + \frac{\beta}{(\bar{\gamma} - \beta)(\gamma - \beta)(\beta - \alpha)} \beta^n + \frac{-\gamma}{(\bar{\gamma} - \gamma)(\gamma - \beta)(\gamma - \alpha)} \gamma^n + \frac{\bar{\gamma}}{(\bar{\gamma} - \gamma)(\bar{\gamma} - \beta)(\bar{\gamma} - \alpha)} \bar{\gamma}^n, \quad n \in \mathbb{N}_0,$$

with

$$\alpha = \frac{1}{4} + \frac{1}{2}\omega + \frac{1}{2}\sqrt{\frac{11}{4} - \omega^2 + \frac{13}{4}\omega^{-1}}, \quad \beta = \frac{1}{4} + \frac{1}{2}\omega - \frac{1}{2}\sqrt{\frac{11}{4} - \omega^2 + \frac{13}{4}\omega^{-1}},$$

$$\gamma = \frac{1}{4} - \frac{1}{2}\omega + \frac{1}{2}\sqrt{\frac{11}{4} - \omega^2 - \frac{13}{4}\omega^{-1}}, \quad \delta = \frac{1}{4} - \frac{1}{2}\omega - \frac{1}{2}\sqrt{\frac{11}{4} - \omega^2 - \frac{13}{4}\omega^{-1}},$$

$$\omega = \sqrt{\frac{11}{12} + \left(\frac{-65}{54} + \sqrt{\frac{563}{108}}\right)^{\frac{1}{3}} + \left(\frac{-65}{54} - \sqrt{\frac{563}{108}}\right)^{\frac{1}{3}}}.$$

The 1-dimensional version of the system (2.2), is the equation

$$x_{n+1} = \frac{x_{n-2}x_{n-1}x_n + x_{n-1}x_{n-2} + x_{n-2} + 1}{x_{n-2}x_{n-1}x_n}, \quad n \in \mathbb{N}_0. \quad (2.3)$$

From the main theorem it follows respectively.

Corollary 2.3. *Let $\{x_n, y_n\}_{n \geq -2}$ be a well defined solution of (2.2). Then, for $n \in \mathbb{N}_0$, one has*

$$x_{2n+1} = \frac{T_{2n+2} + (T_{2n+2} + T_{2n+1})y_{-2} + (T_{2n+4} - T_{2n+3})x_{-1}y_{-2} + T_{2n+3}y_0x_{-1}y_{-2}}{T_{2n+1} + (T_{2n+1} + T_{2n})y_{-2} + (T_{2n+3} - T_{2n+2})x_{-1}y_{-2} + T_{2n+2}y_0x_{-1}y_{-2}},$$

$$x_{2n+2} = \frac{T_{2n+3} + (T_{2n+3} + T_{2n+2})x_{-2} + (T_{2n+5} - T_{2n+4})y_{-1}x_{-2} + T_{2n+4}x_0y_{-1}x_{-2}}{T_{2n+2} + (T_{2n+2} + T_{2n+1})x_{-2} + (T_{2n+4} - T_{2n+3})y_{-1}x_{-2} + T_{2n+3}x_0y_{-1}x_{-2}},$$

$$y_{2n+1} = \frac{T_{2n+2} + (T_{2n+2} + T_{2n+1})x_{-2} + (T_{2n+4} - T_{2n+3})y_{-1}x_{-2} + T_{2n+3}x_0y_{-1}x_{-2}}{T_{2n+1} + (T_{2n+1} + T_{2n})x_{-2} + (T_{2n+3} - T_{2n+2})y_{-1}x_{-2} + T_{2n+2}x_0y_{-1}x_{-2}},$$

$$y_{2n+2} = \frac{T_{2n+3} + (T_{2n+3} + T_{2n+2})y_{-2} + (T_{2n+5} - T_{2n+4})x_{-1}y_{-2} + T_{2n+4}y_0x_{-1}y_{-2}}{T_{2n+2} + (T_{2n+2} + T_{2n+1})y_{-2} + (T_{2n+4} - T_{2n+3})x_{-1}y_{-2} + T_{2n+3}y_0x_{-1}y_{-2}}.$$

Corollary 2.4. Let $\{x_n\}_{n \geq -2}$ be a well defined solution of the equation (2.3). Then for $n \in \mathbb{N}_0$, one has

$$x_{2n+1} = \frac{T_{2n+2} + (T_{2n+2} + T_{2n+1})x_{-2} + (T_{2n+4} - T_{2n+3})x_{-1}x_{-2} + T_{2n+3}x_0x_{-1}x_{-2}}{T_{2n+1} + (T_{2n+1} + T_{2n})x_{-2} + (T_{2n+3} - T_{2n+2})x_{-1}x_{-2} + T_{2n+2}x_0x_{-1}x_{-2}},$$

$$x_{2n+2} = \frac{T_{2n+3} + (T_{2n+3} + T_{2n+2})x_{-2} + (T_{2n+5} - T_{2n+4})x_{-1}x_{-2} + T_{2n+4}x_0x_{-1}x_{-2}}{T_{2n+2} + (T_{2n+2} + T_{2n+1})x_{-2} + (T_{2n+4} - T_{2n+3})x_{-1}x_{-2} + T_{2n+3}x_0x_{-1}x_{-2}}.$$

Remark 2.5. When $a = d = 0$, the system (1.2) takes the form

$$x_{n+1} = \frac{bx_{n-1} + c}{y_n x_{n-1}}, \quad y_{n+1} = \frac{by_{n-1} + c}{x_n y_{n-1}} \quad n \in \mathbb{N}_0. \quad (2.4)$$

As it is noted in [1], the solutions are expressed using Padovan numbers. This system, and some particular cases of it, were the subject of the papers [19,45].

If $d = c = 0$, the system (1.2) becomes

$$x_{n+1} = \frac{ay_n + b}{y_n}, \quad y_{n+1} = \frac{ax_n + b}{x_n}, \quad n \in \mathbb{N}_0. \quad (2.5)$$

Again, it is noted in [1] that:

- The system (2.5) is a particular case of the more general system

$$x_{n+1} = \frac{ay_n + b}{cy_n + d}, \quad y_{n+1} = \frac{\alpha x_n + \beta}{\gamma x_n + \lambda}, \quad n \in \mathbb{N}_0, \quad (2.6)$$

which was completely solved by Stevic in [33] and the solutions are expressed using a generalized Fibonacci sequence.

- Also, particular cases of System (2.6) were studied in [24,28,34,35].

- If also $b = 0$, then the solutions of the system (2.5) are given by

$$\{(x_0, y_0), (a, a), (a, a), \dots\}.$$

3 Proof of the Main Theorem

In order to solve the system (1.2), one needs first to solve the following two homogeneous fourth-order linear difference equations

$$R_{n+1} = aR_n + bR_{n-1} + cR_{n-2} + dR_{n-3}, \quad n \in \mathbb{N}_0, \quad (3.1)$$

$$S_{n+1} = -aS_n + bS_{n-1} - cS_{n-2} + dS_{n-3}, \quad n \in \mathbb{N}_0, \quad (3.2)$$

where the initial values $R_0, R_{-1}, R_{-2}, R_{-3}, S_0, S_{-1}, S_{-2}$ and S_{-3} and the constant coefficients a, b, c and d are real numbers with $d \neq 0$. In fact, one will express the terms of the sequences $(R_n)_{n=-3}^{+\infty}$ and $(S_n)_{n=-3}^{+\infty}$ using the sequence $(J_n)_{n=0}^{+\infty}$.

The difference equation (3.1) has the same characteristic equation as $(J_n)_{n=0}^{+\infty}$, that is the equation (1.6).

To solve the difference equation (3.2) using terms of (3.3), one needs the following fourth-order linear difference equation defined by

$$j_{n+4} = -aj_{n+3} + bj_{n+2} - cj_{n+1} + dj_n, \quad n \in \mathbb{N}_0, \quad (3.3)$$

and the special initial values

$$j_0 = 0, \quad j_1 = 0, \quad j_2 = 1 \text{ and } j_3 = -a. \quad (3.4)$$

The characteristic equation of (3.2) and (3.3) is

$$\lambda^4 + a\lambda^3 - b\lambda^2 + c\lambda - d = 0. \quad (3.5)$$

Clearly the roots of (3.5) are $-\alpha$, $-\beta$, $-\gamma$ and $-\delta$.

Now following the same procedure in solving $\{J_n\}$, it is not hard to see that

$$j_n = (-1)^n J_n.$$

Now, it is possible to prove the following result.

Lemma 3.1. *One has for all $n \in \mathbb{N}_0$,*

$$R_n = dJ_{n+1}R_{-3} + (cJ_{n+1} + dJ_n)R_{-2} + (J_{n+3} - aJ_{n+2})R_{-1} + J_{n+2}R_0, \quad (3.6)$$

$$S_n = (-1)^{n+1} [dJ_{n+1}S_{-3} - (cJ_{n+1} + dJ_n)S_{-2} + (J_{n+3} - aJ_{n+2})S_{-1} - J_{n+2}S_0]. \quad (3.7)$$

Proof. Assume that α , β , γ and δ are the distinct roots of the characteristic equation (1.6), so

$$R_n = c'_1\alpha^n + c'_2\beta^n + c'_3\gamma^n + c'_4\delta^n, \quad n \geq -3.$$

Using the initial values R_0, R_{-1}, R_{-2} and R_{-3} , one get

$$\begin{cases} \frac{1}{\alpha^3}c'_1 + \frac{1}{\beta^3}c'_2 + \frac{1}{\gamma^3}c'_3 + \frac{1}{\delta^3}c'_4 & = R_{-3} \\ \frac{1}{\alpha^2}c'_1 + \frac{1}{\beta^2}c'_2 + \frac{1}{\gamma^2}c'_3 + \frac{1}{\delta^2}c'_4 & = R_{-2} \\ \frac{1}{\alpha}c'_1 + \frac{1}{\beta}c'_2 + \frac{1}{\gamma}c'_3 + \frac{1}{\delta}c'_4 & = R_{-1} \\ c'_1 + c'_2 + c'_3 + c'_4 & = R_0, \end{cases} \quad (3.8)$$

after some calculations using the Cramer method one get

$$\begin{aligned}
 c'_1 &= \frac{\beta\gamma\delta\alpha^3}{(\delta-\alpha)(\gamma-\alpha)(\beta-\alpha)}R_{-3} - \frac{(\gamma\beta+\gamma\delta+\beta\delta)\alpha^3}{(\delta-\alpha)(\gamma-\alpha)(\beta-\alpha)}R_{-2} \\
 &\quad + \frac{(\beta+\gamma+\delta)\alpha^3}{(\delta-\alpha)(\gamma-\alpha)(\beta-\alpha)}R_{-1} - \frac{\alpha^3}{(\delta-\alpha)(\gamma-\alpha)(\beta-\alpha)}R_0 \\
 c'_2 &= -\frac{\alpha\gamma\delta\beta^3}{(\delta-\beta)(\gamma-\beta)(\beta-\alpha)}R_{-3} + \frac{(\gamma\alpha+\gamma\delta+\alpha\delta)\beta^3}{(\delta-\beta)(\gamma-\beta)(\beta-\alpha)}R_{-2} \\
 &\quad - \frac{(\alpha+\gamma+\delta)\beta^3}{(\delta-\beta)(\gamma-\beta)(\beta-\alpha)}R_{-1} + \frac{\beta^3}{(\delta-\beta)(\gamma-\beta)(\beta-\alpha)}R_0 \\
 c'_3 &= \frac{\alpha\beta\delta\gamma^3}{(\delta-\gamma)(\gamma-\beta)(\gamma-\alpha)}R_{-3} - \frac{(\alpha\beta+\alpha\delta+\beta\delta)\gamma^3}{(\delta-\gamma)(\gamma-\beta)(\gamma-\alpha)}R_{-2} \\
 &\quad + \frac{(\alpha+\beta+\delta)\gamma^3}{(\delta-\gamma)(\gamma-\beta)(\gamma-\alpha)}R_{-1} - \frac{\gamma^3}{(\delta-\gamma)(\gamma-\beta)(\gamma-\alpha)}R_0 \\
 c'_4 &= -\frac{\alpha\beta\gamma\delta^3}{(\delta-\gamma)(\delta-\beta)(\delta-\alpha)}R_{-3} + \frac{(\gamma\alpha+\gamma\beta+\alpha\beta)\delta^3}{(\delta-\gamma)(\delta-\beta)(\delta-\alpha)}R_{-2} \\
 &\quad - \frac{(\alpha+\beta+\gamma)\delta^3}{(\delta-\gamma)(\delta-\beta)(\delta-\alpha)}R_{-1} + \frac{\delta^3}{(\delta-\gamma)(\delta-\beta)(\delta-\alpha)}R_0
 \end{aligned}$$

that is,

$$\begin{aligned}
 R_n &= \left(\frac{\beta\gamma\delta\alpha^3}{(\delta-\alpha)(\gamma-\alpha)(\beta-\alpha)}\alpha^n - \frac{\alpha\gamma\delta\beta^3}{(\delta-\beta)(\gamma-\beta)(\beta-\alpha)}\beta^n + \frac{\alpha\beta\delta\gamma^3}{(\delta-\gamma)(\gamma-\beta)(\gamma-\alpha)}\gamma^n \right. \\
 &\quad \left. - \frac{\alpha\beta\gamma\delta^3}{(\delta-\gamma)(\delta-\beta)(\delta-\alpha)}\delta^n \right) R_{-3} \\
 &\quad + \left(-\frac{(\gamma\beta+\gamma\delta+\beta\delta)\alpha^3}{(\delta-\alpha)(\gamma-\alpha)(\beta-\alpha)}\alpha^n + \frac{(\gamma\alpha+\gamma\delta+\alpha\delta)\beta^3}{(\delta-\beta)(\gamma-\beta)(\beta-\alpha)}\beta^n - \frac{(\alpha\beta+\alpha\delta+\beta\delta)\gamma^3}{(\delta-\gamma)(\gamma-\beta)(\gamma-\alpha)}\gamma^n \right. \\
 &\quad \left. + \frac{(\gamma\alpha+\gamma\beta+\alpha\beta)\delta^3}{(\delta-\gamma)(\delta-\beta)(\delta-\alpha)}\delta^n \right) R_{-2} \\
 &\quad + \left(\frac{(\beta+\gamma+\delta)\alpha^3}{(\delta-\alpha)(\gamma-\alpha)(\beta-\alpha)}\alpha^n - \frac{(\alpha+\gamma+\delta)\beta^3}{(\delta-\beta)(\gamma-\beta)(\beta-\alpha)}\beta^n + \frac{(\alpha+\beta+\delta)\gamma^3}{(\delta-\gamma)(\gamma-\beta)(\gamma-\alpha)}\gamma^n \right. \\
 &\quad \left. - \frac{(\alpha+\beta+\gamma)\delta^3}{(\delta-\gamma)(\delta-\beta)(\delta-\alpha)}\delta^n \right) R_{-1} \\
 &\quad + \left(-\frac{\alpha^3}{(\delta-\alpha)(\gamma-\alpha)(\beta-\alpha)}\alpha^n + \frac{\beta^3}{(\delta-\beta)(\gamma-\beta)(\beta-\alpha)}\beta^n - \frac{\gamma^3}{(\delta-\gamma)(\gamma-\beta)(\gamma-\alpha)}\gamma^n \right. \\
 &\quad \left. + \frac{\delta^3}{(\delta-\gamma)(\delta-\beta)(\delta-\alpha)}\delta^n \right) R_0.
 \end{aligned}$$

$$R_n = dJ_{n+1}R_{-3} + (cJ_{n+1} + dJ_n)R_{-2} + (J_{n+3} - aJ_{n+2})R_{-1} + J_{n+2}R_0.$$

The proof of the other cases is similar and will be omitted.

Let $A := -a$, $B := b$, $C := -c$ and $D := d$ then, equation (3.2) takes the form (3.1) and the equation (3.3) takes the form (3.3). Then analogous to the formula of (3.1) one obtains

$$S_n = Dj_{n+1}S_{-3} + (Cj_{n+1} + Dj_n)S_{-2} + (j_{n+3} - Aj_{n+2})S_{-1} + j_{n+2}S_0.$$

Using the fact that $j_n = (-1)^n J_n$, $A = -a$ and $C := -c$ one get

$$S_n = (-1)^{n+1} [dJ_{n+1}S_{-3} - (cJ_{n+1} + dJ_n)S_{-2} + (J_{n+3} - aJ_{n+2})S_{-1} - J_{n+2}S_0].$$

□

Proof of the Main Theorem.

Replacing

$$x_n = \frac{u_n}{v_{n-1}}, \quad y_n = \frac{v_n}{u_{n-1}}, \quad n \geq -2, \quad (3.9)$$

in system (1.2) one get the following linear fourth-order system of difference equations

$$u_{n+1} = av_n + bu_{n-1} + cv_{n-2} + du_{n-3}, \quad v_{n+1} = au_n + bv_{n-1} + cu_{n-2} + dv_{n-3}, \quad n \in \mathbb{N}_0, \quad (3.10)$$

where the initial values $u_{-3}, u_{-2}, u_{-1}, u_0, v_{-3}, v_{-2}, v_{-1}, v_0$ are nonzero real numbers.From (3.10) one has for $n \in \mathbb{N}_0$,

$$\begin{cases} u_{n+1} + v_{n+1} = a(v_n + u_n) + b(u_{n-1} + v_{n-1}) + c(v_{n-2} + u_{n-2}) + d(u_{n-3} + v_{n-3}), \\ u_{n+1} - v_{n+1} = a(v_n - u_n) + b(u_{n-1} - v_{n-1}) + c(v_{n-2} - u_{n-2}) + d(u_{n-3} - v_{n-3}). \end{cases}$$

Putting again

$$R_n = u_n + v_n, \quad S_n = u_n - v_n, \quad n \geq -3, \quad (3.11)$$

one obtains two fourth-order homogeneous linear difference equations:

$$R_{n+1} = aR_n + bR_{n-1} + cR_{n-2} + dR_{n-3}, \quad n \in \mathbb{N}_0,$$

and

$$S_{n+1} = -aS_n + bS_{n-1} - cS_{n-2} + dS_{n-3}, \quad n \in \mathbb{N}_0. \quad (3.12)$$

Using (3.11), one get for $n \geq -3$,

$$u_n = \frac{1}{2}(R_n + S_n), \quad v_n = \frac{1}{2}(R_n - S_n).$$

From Lemma 3.1 one obtains,

$$\begin{cases} u_{2n-1} = \frac{1}{2} [dJ_{2n}(R_{-3} + S_{-3}) + (cJ_{2n} + dJ_{2n-1})(R_{-2} - S_{-2}) + (J_{2n+2} - aJ_{2n+1})(R_{-1} + S_{-1}) \\ \quad + J_{2n+1}(R_0 - S_0)], \quad n \in \mathbb{N}, \\ u_{2n} = \frac{1}{2} [dJ_{2n+1}(R_{-3} - S_{-3}) + (cJ_{2n+1} + dJ_{2n})(R_{-2} + S_{-2}) + (J_{2n+3} - aJ_{2n+2})(R_{-1} - S_{-1}) \\ \quad + J_{2n+2}(R_0 + S_0)], \quad n \in \mathbb{N}_0, \end{cases} \quad (3.13)$$

$$\begin{cases} v_{2n-1} = \frac{1}{2} [dJ_{2n}(R_{-3} - S_{-3}) + (cJ_{2n} + dJ_{2n-1})(R_{-2} + S_{-2}) + (J_{2n+2} - aJ_{2n+1})(R_{-1} - S_{-1}) \\ \quad + J_{2n+1}(R_0 + S_0)], \quad n \in \mathbb{N}, \\ v_{2n} = \frac{1}{2} [dJ_{2n+1}(R_{-3} + S_{-3}) + (cJ_{2n+1} + dJ_{2n})(R_{-2} - S_{-2}) + (J_{2n+3} - aJ_{2n+2})(R_{-1} + S_{-1}) \\ \quad + J_{2n+2}(R_0 - S_0)], \quad n \in \mathbb{N}_0. \end{cases} \quad (3.14)$$

Substituting (3.13) and (3.14) in (3.9), one get for $n \in \mathbb{N}_0$,

$$x_{2n+1} = \frac{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1}) \frac{R_{-2} - S_{-2}}{R_{-3} + S_{-3}} + (J_{2n+4} - aJ_{2n+3}) \frac{R_{-1} + S_{-1}}{R_{-3} + S_{-3}} + J_{2n+3} \frac{R_0 - S_0}{R_{-3} + S_{-3}}}{dJ_{2n+1} + (cJ_{2n+1} + dJ_{2n}) \frac{R_{-2} - S_{-2}}{R_{-3} + S_{-3}} + (J_{2n+3} - aJ_{2n+2}) \frac{R_{-1} + S_{-1}}{R_{-3} + S_{-3}} + J_{2n+2} \frac{R_0 - S_0}{R_{-3} + S_{-3}}}, \quad (3.15)$$

$$x_{2n+2} = \frac{dJ_{2n+3} + (cJ_{2n+3} + dJ_{2n+2}) \frac{R_{-2} + S_{-2}}{R_{-3} - S_{-3}} + (J_{2n+5} - aJ_{2n+4}) \frac{R_{-1} - S_{-1}}{R_{-3} - S_{-3}} + J_{2n+4} \frac{R_0 + S_0}{R_{-3} - S_{-3}}}{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1}) \frac{R_{-2} + S_{-2}}{R_{-3} - S_{-3}} + (J_{2n+4} - aJ_{2n+3}) \frac{R_{-1} - S_{-1}}{R_{-3} - S_{-3}} + J_{2n+3} \frac{R_0 + S_0}{R_{-3} - S_{-3}}}, \quad (3.16)$$

$$y_{2n+1} = \frac{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1}) \frac{R_{-2} + S_{-2}}{R_{-3} - S_{-3}} + (J_{2n+4} - aJ_{2n+3}) \frac{R_{-1} - S_{-1}}{R_{-3} - S_{-3}} + J_{2n+3} \frac{R_0 + S_0}{R_{-3} - S_{-3}}}{dJ_{2n+1} + (cJ_{2n+1} + dJ_{2n}) \frac{R_{-2} + S_{-2}}{R_{-3} - S_{-3}} + (J_{2n+3} - aJ_{2n+2}) \frac{R_{-1} - S_{-1}}{R_{-3} - S_{-3}} + J_{2n+2} \frac{R_0 + S_0}{R_{-3} - S_{-3}}}, \quad (3.17)$$

and

$$y_{2n+2} = \frac{dJ_{2n+3} + (cJ_{2n+3} + dJ_{2n+2}) \frac{R_{-2} - S_{-2}}{R_{-3} + S_{-3}} + (J_{2n+5} - aJ_{2n+4}) \frac{R_{-1} + S_{-1}}{R_{-3} + S_{-3}} + J_{2n+4} \frac{R_0 - S_0}{R_{-3} + S_{-3}}}{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1}) \frac{R_{-2} - S_{-2}}{R_{-3} + S_{-3}} + (J_{2n+4} - aJ_{2n+3}) \frac{R_{-1} + S_{-1}}{R_{-3} + S_{-3}} + J_{2n+3} \frac{R_0 - S_0}{R_{-3} + S_{-3}}}. \quad (3.18)$$

One has

$$x_{-2} = \frac{u_{-2}}{v_{-3}} = \frac{R_{-2} + S_{-2}}{R_{-3} - S_{-3}}, \quad x_{-1} = \frac{u_{-1}}{v_{-2}} = \frac{R_{-1} + S_{-1}}{R_{-2} - S_{-2}}, \quad x_0 = \frac{u_0}{v_{-1}} = \frac{R_0 + S_0}{R_{-1} - S_{-1}}, \quad (3.19)$$

$$y_{-2} = \frac{v_{-2}}{u_{-3}} = \frac{R_{-2} - S_{-2}}{R_{-3} + S_{-3}}, \quad y_{-1} = \frac{v_{-1}}{u_{-2}} = \frac{R_{-1} - S_{-1}}{R_{-2} + S_{-2}}, \quad y_0 = \frac{v_0}{u_{-1}} = \frac{R_0 - S_0}{R_{-1} + S_{-1}}. \quad (3.20)$$

From (3.19), (3.20) one get,

$$\begin{cases} \frac{R_{-1} + S_{-1}}{R_{-3} + S_{-3}} = \frac{R_{-1} + S_{-1}}{R_{-2} - S_{-2}} \times \frac{R_{-2} - S_{-2}}{R_{-3} + S_{-3}} = x_{-1}y_{-2}, \\ \frac{R_0 - S_0}{R_{-3} + S_{-3}} = \frac{R_0 - S_0}{R_{-1} + S_{-1}} \times \frac{R_{-1} + S_{-1}}{R_{-2} - S_{-2}} \times \frac{R_{-2} - S_{-2}}{R_{-3} + S_{-3}} = y_0x_{-1}y_{-2}, \end{cases} \quad (3.21)$$

$$\begin{cases} \frac{R_{-1} - S_{-1}}{R_{-3} - S_{-3}} = \frac{R_{-1} - S_{-1}}{R_{-2} + S_{-2}} \times \frac{R_{-2} + S_{-2}}{R_{-3} - S_{-3}} = y_{-1}x_{-2}, \\ \frac{R_0 + S_0}{R_{-3} - S_{-3}} = \frac{R_0 + S_0}{R_{-1} - S_{-1}} \times \frac{R_{-1} - S_{-1}}{R_{-2} + S_{-2}} \times \frac{R_{-2} + S_{-2}}{R_{-3} - S_{-3}} = x_0y_{-1}x_{-2}. \end{cases} \quad (3.22)$$

Using (3.15), (3.16), (3.17), (3.18), (3.21) and (3.22), one obtains the closed form of the solutions of the system (1.2), that is for $n \in \mathbb{N}_0$, one has

$$\begin{cases} x_{2n+1} = \frac{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1})y_{-2} + (J_{2n+4} - aJ_{2n+3})x_{-1}y_{-2} + J_{2n+3}y_0x_{-1}y_{-2}}{dJ_{2n+1} + (cJ_{2n+1} + dJ_{2n})y_{-2} + (J_{2n+3} - aJ_{2n+2})x_{-1}y_{-2} + J_{2n+2}y_0x_{-1}y_{-2}}, \\ x_{2n+2} = \frac{dJ_{2n+3} + (cJ_{2n+3} + dJ_{2n+2})x_{-2} + (J_{2n+5} - aJ_{2n+4})y_{-1}x_{-2} + J_{2n+4}x_0y_{-1}x_{-2}}{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1})x_{-2} + (J_{2n+4} - aJ_{2n+3})y_{-1}x_{-2} + J_{2n+3}x_0y_{-1}x_{-2}}, \end{cases}$$

$$\begin{cases} y_{2n+1} = \frac{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1})x_{-2} + (J_{2n+4} - aJ_{2n+3})y_{-1}x_{-2} + J_{2n+3}x_0y_{-1}x_{-2}}{dJ_{2n+1} + (cJ_{2n+1} + dJ_{2n})x_{-2} + (J_{2n+3} - aJ_{2n+2})y_{-1}x_{-2} + J_{2n+2}x_0y_{-1}x_{-2}}, \\ y_{2n+2} = \frac{dJ_{2n+3} + (cJ_{2n+3} + dJ_{2n+2})y_{-2} + (J_{2n+5} - aJ_{2n+4})x_{-1}y_{-2} + J_{2n+4}y_0x_{-1}y_{-2}}{dJ_{2n+2} + (cJ_{2n+2} + dJ_{2n+1})y_{-2} + (J_{2n+4} - aJ_{2n+3})x_{-1}y_{-2} + J_{2n+3}y_0x_{-1}y_{-2}}. \end{cases}$$

Remark 3.2. - The content of the present paper was posted on arXiv on 31.10.2019, ref. arXiv:1910.14365.

- Some parts of the results of this paper were used in the reference [2] in which the authors have generalized the system (1.2).

Acknowledgements

The authors would like to thank the two referees for their comments and suggestions. Also, the authors acknowledge the support of the Algerian General Directorate for Scientific Research and Technological Development DGRSDT.

Conflict of Interest

The authors have no conflicts of interest to declare.

References

- [1] Y. AKROUR, N. TOUAFEK AND Y. HALIM, *On a system of difference equations of second order solved in closed form*, Miskolc Math. Notes **20**(2) (2019), 701–717.
- [2] Y. AKROUR, M. KARA, N. TOUAFEK AND Y. YAZLIK, *Solutions formulas for some general systems of difference equations*, Miskolc Math. Notes in press (2021).
- [3] A. ASIRI, M. M. EL-DESSOKY AND E. M. ELSAYED, *Solution of a third order fractional system of difference equations*, J. Comput. Anal. Appl. **24**(3) (2018), 444–453.
- [4] R. AZIZI, *Global behaviour of the rational Riccati difference equation of order two: the general case*, J. Difference Equ. Appl. **18**(6) (2012), 947–961.
- [5] F. BELHANNACHE, N. TOUAFEK, AND R. ABO-ZEID, *Dynamics of a third-order rational difference equation*, Bull. Math. Soc. Sci. Math. Roumanie (N.S.) **59**(107) (2016), 13–22.
- [6] M. M. EL-DESSOKY, *On a solvable for some systems of rational difference equations*, J. Nonlinear Sci. Appl. **9**(6) (2016), 3744–3759.
- [7] M. M. EL-DESSOKY, E. M. ELSAYED, E. M. ELABBASY AND ASIM ASIRI, *Expressions of the solutions of some systems of difference equations*, J. Comput. Anal. Appl. **27**(7) (2019), 1161–1172.
- [8] E. M. ELSAYED, *On a system of two nonlinear difference equations of order two*, Proc. Jangjeon Math. Soc. **18**(1) (2015), 353–368.
- [9] E. M. ELSAYED AND T. F. IBRAHIM, *Periodicity and solutions for some systems of nonlinear rational difference equations*, Hacet. J. Math. Stat. **44**(1) (2015), 1361–1390.
- [10] E. M. ELSAYED, *Solution for systems of difference equations of rational form of order two*, Comput. Appl. Math. **33**(1) (2015), 751–765.

- [11] M. GUMUS, *The global asymptotic stability of a system of difference equations*, J. Difference Equ. Appl. **24**(1) (2018), 976–991.
- [12] M. GUMUS AND O. OCALAN, *The qualitative analysis of a rational system of difference equations*, J. Fract. Calc. Appl. **9**(2) (2018), 113–126.
- [13] M. GUMUS AND R. ABO-ZEID, *On the solutions of a $(2k+2)$ th order difference equation*, Dyn. Contin. Discrete Impuls. Syst. Ser. B Appl. Algorithms **25**(1) (2018), 129–143.
- [14] M. GUMUS, *Analysis of periodicity for a new class of nonlinear difference equations by using a new method*, Electron. J. Math. Anal. Appl. **8**(1) (2020), 109–116.
- [15] M. GUMUS, *The periodic character in a higher order difference equation with delays*, Math. Methods Appl. Sci. **43**(3) (2020), 1112–1123.
- [16] N. HADDAD, N. TOUAFEK AND J. F. T. RABAGO, *Solution form of a higher-order system of difference equations and dynamical behavior of its special case*, Math. Methods Appl. Sci. **40**(1) (2017), 3599–3607.
- [17] N. HADDAD, N. TOUAFEK AND E. M. ELSAYED, *A note on a system of difference equations*, An. Ştiinţ. Univ. Al. I. Cuza Iaşi, Ser. Nouă, Mat.(N.S.) **63**(3) (2017), 599–606.
- [18] N. HADDAD, N. TOUAFEK AND J. F. T. RABAGO, *Well-defined solutions of a system of difference equations*, J. Appl. Math. Comput. **56**(1) (2018), 439–458.
- [19] Y. HALIM AND J. F. T. RABAGO, *On the solutions of a second-order difference equation in terms of generalized Padovan sequences*, Math. Slovaca. **68**(3) (2018), 625–638.
- [20] Y. HALIM AND J. F. T. RABAGO, *On some solvable systems of difference equations with solutions associated to Fibonacci numbers*, Electron. J. Math. Anal. Appl. **5**(1) (2017), 166–178.
- [21] Y. HALIM, *A system of difference equations with solutions associated to Fibonacci numbers*, Int. J. Differ. Equ. **11**(1) (2016), 65–77.
- [22] Y. HALIM AND M. BAYRAM, *On the solutions of a higher-order difference equation in terms of generalized Fibonacci sequences*, Math. Methods Appl. Sci. **39**(1) (2016), 2974–2982.
- [23] Y. HALIM, N. TOUAFEK AND E. M. ELSAYED, *Closed form solution of some systems of rational difference equations in terms of Fibonacci numbers*, Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal. **21**(6) (2014), 473–486.
- [24] Y. HALIM, *Global character of systems of rational difference equations*, Electron. J. Math. Anal. Appl. **3**(1) (2016), 204–214.
- [25] G. S. HATHIWALA AND D. V. SHAH, *Binet-type formula for the sequence of Tetranacci numbers by alternate methods*, J. Interdiscip. Math. **6**(1) (2017), 37–48.
- [26] M. KARA AND Y. YAZLIK, *Solvability of a system of nonlinear difference equations of higher order*, Turkish J. Math. **43**(3) (2019), 1533–1565.
- [27] M. KARA AND Y. YAZLIK, *Solvability of a nonlinear three-dimensional system of difference equations with constant coefficients*, Math. Slovaca. **71**(5) (2021), 1133–1148.

- [28] H. MATSUNAGA AND R. SUZUKI, *Classification of global behavior of a system of rational difference equations*, Appl. Math. Lett. **85**(1) (2018), 57–63.
- [29] S. STEVIC, *Some representations of the general solution to a difference equation of additive type*, Adv. Differ. Equ. **2019**(431) (2019), 1–19.
- [30] S. STEVIC, B. IRICANIN AND W. KOSMALA, *Representations of general solutions to some classes of nonlinear difference equations*, Adv. Differ. Equ. **2019** (73) (2019), 1–21.
- [31] S. STEVIC, B. IRICANIN, W. KOSMALA AND Z. SMARDA, *Note on the bilinear difference equation with a delay*, Math. Methods Appl. Sci. **41**(18) (2018), 9349–9360.
- [32] S. STEVIC, B. IRICANIN, W. KOSMALA AND Z. SMARDA, *Representation of solutions of a solvable nonlinear difference equation of second order*, Electron. J. Qual. theory Differ. Equ. **2015** (95) (2018), 1–18.
- [33] S. STEVIC, *Representation of solutions of bilinear difference equations in terms of generalized Fibonacci sequences*, Electron. J. Qual. theory Differ. Equ. **2014**(67) (2014), 1–15.
- [34] D. T. TOLLU, Y. YAZLIK AND N. TASKARA, *On the solutions of two special types of Riccati difference equation via Fibonacci numbers*, Adv. Differ. Equ. **2013**(174) (2013), 1–7.
- [35] D. T. TOLLU, Y. YAZLIK AND N. TASKARA, *The solutions of four Riccati difference equations associated with Fibonacci numbers*, Balkan J. Math. **2**(1) (2014), 163–172.
- [36] D. T. TOLLU, Y. YAZLIK AND N. TASKARA, *On fourteen solvable systems of difference equations*, Appl. Math. Comput. **233**(1) (2014), 310–319.
- [37] N. TOUAFEK, *On some fractional systems of difference equations*, Iran. J. Math. Sci. Inform. **9**(2) (2014), 303–305.
- [38] N. TOUAFEK, Y. HALIM, *On max type difference equations: expressions of solutions*, Int. j. appl. nonlinear sci. **11**(4) (2011), 396–402.
- [39] N. TOUAFEK, E.M. ELSAYED, *On the periodicity of some systems of nonlinear difference equations*, Bull. math. Soc. Sci. Math. Roum. **55**(1) (2012), 217–224.
- [40] N. TOUAFEK, E.M. ELSAYED, *On the solutions of systems of rational difference equations*, Math. Comput. Model. **55**(1) (2012), 1987–1997.
- [41] C. WANG, X. FANG AND R. LI, *On the solution for a system of two rational difference equations*, J. Comput. Anal. Appl. **20**(1) (2016), 175–186.
- [42] C. WANG, X. FANG AND R. LI, *On the dynamics of a certain four-order fractional difference equations*, J. Comput. Anal. Appl. **22**(5) (2017), 968–976.
- [43] C. WANG, X. FANG AND R. LI, *On the periodicity of a max-type rational difference equation*, J. Nonlinear Sci. Appl. **10**(9) (2017), 4648–4661.
- [44] C. WANG, Y. ZHOU, S. PAN, R. LI, *On a system of three max-type nonlinear difference equations*, J. Comput. Anal. Appl. **25**(8) (2018), 1463–1479.
- [45] Y. YAZLIK, D. T. TOLLU, N. TASKARA, *On the solutions of difference equation systems with Padovan numbers*, Appl. Math. **4** (2013), 15–20.

- [46] Y. YAZLIK, M. KARA, *On a solvable system of difference equations of higher-order with period two coefficients*, Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat. **68**(1) (2019), 1675–1693.
- [47] M. E. WADDILL, *The Tetranacci sequence and generalizations*, Fibonacci Quart. **30**(1) (1992), 9–20.